

1) The ground state of the hydrogen atom and the uncertainty principle

Bohr : $E_n = -\frac{me^4}{2\hbar^2} \frac{1}{n^2}$
 $n=1, 2, 3, \dots$

"Why" is no energy lower than $E_1 = -\frac{me^4}{2\hbar^2}$ possible?

$$E = \frac{p^2}{2m} - \frac{e^2}{r}, \quad \Delta p \Delta r \gtrsim \hbar$$

Energy is lowered by decreasing r and/or p . Best we can

do is

2

$$\bullet E \sim \frac{\Delta p^2}{2m} - \frac{e^2}{\Delta r}$$

$$= \frac{\Delta p^2}{2m} - \frac{e^2 \Delta p}{\hbar} \quad \left(\text{using } \Delta r = \frac{\hbar}{\Delta p} \right)$$

$$\text{min } E \Rightarrow 0 = \frac{\partial E}{\partial \Delta p} = \frac{\Delta p}{m} - \frac{e^2}{\hbar}$$

$$\bullet \Delta p = \frac{me^2}{\hbar}$$

$$\Delta r = \frac{\hbar}{\Delta p} = \frac{\hbar^2}{me^2}$$

(Bohr radius!)

$$E = \frac{1}{2m} \left(\frac{me^2}{\hbar} \right)^2 - \frac{e^2 me^2}{\hbar^2} = -\frac{1}{2} \frac{me^4}{\hbar^2} = -13.6 \text{ eV}$$


Thus the first Bohr orbital

is the lowest possible energy

- of the hydrogen atom, 3
- consistent with the uncertainty principle!

2) Wave-particle duality:

The double-slit experiment and the uncertainty principle




- An electron gun emits electrons of energy E_0 toward a screen with

two slits, with separation d . (4

What is observed on a screen a distance $L \gg d$ away?

Electrons are observed to hit the screen one by one at particular points x , corresponding to angles $\theta \approx x/L$. After

a long time, if we count all the electrons which have hit the screen, we find



The maximum numbers of counts occur for constructive interference of the de Broglie waves :

$$n = 0, \pm 1, \pm 2, \dots$$

$$d \sin \theta = n \lambda_e, \quad \lambda_e = \frac{h}{p_0}$$

The minima correspond to destructive interference :

$$d \sin \theta = \left(n + \frac{1}{2}\right) \lambda_e, \quad n = 0, \pm 1, \pm 2,$$

Nonetheless, each electron hits the screen in one and only one place. How can we reconcile the interference pattern with the very

- discrete nature of the (6
● electron?

Proposition A: Each electron passes through either slit 1 or slit 2.

- The particle nature of the electron would seem to imply the validity of Prop. A. Yet the interference pattern accumulates even if we slow down the electron source so that at most one electron impinges on the two slits at any given time. Interference would seem to require the
-

- electron to pass through both slits. 7

To resolve this apparent paradox, we should test prop. A directly, e.g., by shining some light on the slits, to see

- which slit the electron passes through. In order to resolve which slit the electron went

through, we need to use light of wavelength $\lambda \leq d$.

If the light source is bright enough, then each

- electron scatters one or more photons in passing

- through the slits, and we find that prop. A is indeed true. However, each photon has momentum

$p_y = \frac{h}{\lambda_y}$. The momentum

- p_0 of the electron is thus changed by an amount

$\Delta p \sim \frac{h}{\lambda_y}$, which changes

the "trajectory" by an amount $\Delta \theta \sim \frac{\Delta p}{p_0} = \frac{\lambda_e}{\lambda_y}$.

- The angular separation of

- neighboring maxima and minima of the interference pattern is 9

$$d |\sin \theta_{\max} - \sin \theta_{\min}| = \frac{\lambda e}{2}$$

$$|\theta_{\max} - \theta_{\min}| \sim \frac{\lambda e}{2d}$$

- If $\Delta \theta > \frac{\lambda e}{2d}$, then the interference pattern will be smeared out, i.e., if

$$\Delta \theta \sim \frac{\lambda e}{\lambda y} > \frac{\lambda e}{2d} \Rightarrow \lambda y < 2d.$$

- But it is necessary to use light with $\lambda y < d$ to resolve which slit the

- electron passed through! (10)
Thus, if we check prop. A,
the interference pattern
is destroyed. The

Seemingly contradictory

wave-like and particle-like

aspects of the electron
cannot be brought into

conflict, because observation

of one aspect precludes

observation of the other

aspect. This was shown

for the specific example

of light-scattering, but 11
Heisenberg postulated that
this is a fundamental
feature of the quantum
world: It is impossible

to design any apparatus
capable of simultaneously

determining the position
and momentum of an object
with a joint precision
violating the inequality

$$\Delta x \Delta p_x \leq \frac{\hbar}{2}.$$