

# Physics 371 lecture 31

## Magnetic moment

Definition (classical)

$$\vec{\mu} = \frac{1}{2c} \int d^3r (\vec{r} \times \vec{j}_g(\vec{r}))$$

$$\bullet QM: \quad \vec{j}_g = \frac{e}{2M} \left( \psi^* \frac{\hbar}{i} \nabla \psi - \psi \frac{\hbar}{i} \nabla \psi^* \right) \\ = \frac{e}{M} \operatorname{Re} [\psi^* \vec{p} \psi]$$

$$\begin{aligned} \langle \vec{\mu} \rangle &= \frac{e}{2Mc} \int d^3r (\vec{r} \times \operatorname{Re} [\psi^* \vec{p} \psi]) \\ &= \frac{e}{2Mc} \operatorname{Re} \int d^3r \psi^* (\vec{r} \times \vec{p}) \psi \\ &= \frac{e}{2Mc} \langle \vec{l} \rangle \end{aligned}$$

## Magnetic moment operator

$$\vec{\mu} = \frac{g}{2mc} \vec{L}$$

## Zeeman effect

Classically,  $E = -\vec{\mu} \cdot \vec{B}$

$$\Rightarrow \hat{H} = -\vec{\mu} \cdot \vec{B}$$

Say  $\vec{B} = B \hat{z}$  (constant field)

$$\hat{H} = -\frac{gB}{2mc} \hat{L}_z$$

$$\hat{H}|Y_{lm}\rangle = -\frac{g\hbar m}{2mc} B |Y_{lm}\rangle$$

↑ magnetic quantum #

$$E_m = -\frac{g\hbar B}{2mc} m, \quad m = -l, \dots, l$$

$$\Delta t \geq \frac{\hbar/2}{g\hbar B/\hbar Mc} = \frac{Mc}{gB}$$

Classically, a magnetic field exerts torque on a magnetic moment:

$$\vec{\tau} = \vec{\mu} \times \vec{B} = \frac{d\vec{L}}{dt}$$

Quantum mechanically, this equation of motion still holds on average:

$$\frac{d}{dt} \langle \vec{L} \rangle = \langle \vec{\mu} \rangle \times \vec{B}$$

Hw: Prove this!


$$\frac{d}{dt} \langle \vec{L} \rangle = -\frac{g\vec{B}}{2Mc} \times \langle \vec{L} \rangle = \vec{\Omega} \times \langle \vec{L} \rangle$$

The average angular momentum  
vector precesses about the  
magnetic field with  
angular frequency

$$\Omega = |\vec{\omega}| = \frac{gB}{2mc}$$

Larmor frequency

$$\Omega \Delta t \geq \frac{gB}{2mc} \frac{mc}{gB} = \frac{1}{2}$$



$$\frac{\Delta L_{\perp}}{L_{\perp}} \sim \frac{1}{2}$$

Thus, applying a magnetic field in the  $z$ -direction for a time  $\Delta t$  allows one to resolve the different Zeeman split levels, and hence to measure the  $z$ -component of the angular momentum. However, such a measurement necessarily perturbs  $L_x$  and  $L_y$ , consistent with the uncertainty relation

$$\Delta L_x \Delta L_y \geq \frac{\hbar}{2} |\langle L_z \rangle|.$$

## Intrinsic magnetic moment

[7]

There is also a magnetic moment associated with the spin of the electron. However, because  $\vec{S} \neq \vec{r} \times \vec{p}$ , the derivation  $\vec{\mu} = -\frac{e}{2mc} \vec{S}$  does not work. Instead,

$$\vec{\mu} = g \left( -\frac{e}{2mc} \right) \vec{S},$$

where  $g \approx 2$  (this factor comes from relativistic quantum mechanics, and is one of the most precisely known quantities in the physical sciences).

The Zeeman effect for the spin

is thus (for  $\vec{B} = B \hat{z} = \text{const.}$ ) (8)


$$-\vec{\mu} \cdot \vec{B} = \frac{geB}{2mc} S_z = \pm \frac{g}{2} \frac{e\hbar}{2mc} B$$
$$= \pm \frac{g}{2} \mu_B B$$

### Stern-Gerlach experiment

In an inhomogeneous magnetic field, there is not only a torque, but also a force on a magnetic moment

$$\vec{F} = D(\vec{\mu} \cdot \vec{B}).$$

This force can be used to separate out the different  $m$ -components of a beam of particles



For an unpolarized beam of particles, the # of components in the beam downstream of the magnet determines the total angular momentum quantum #:  $N = 2l + 1$ .


For an individual particle, the deflection represents a measurement (determination) of the z-component of the angular momentum.

Let's focus on the simplest case,  $s=1/2$ . It would be difficult to do the experiment with a free electron, because the beam


would be bent and dispersed by [10] the magnetic field, due to the Lorentz force. But for a neutral particle, such as a hydrogen atom, the charge of the electron is balanced by that of the proton, but the magnetic moment is not since

$$\mu_B = \frac{e\hbar}{2m_e c} \gg \mu_p \sim \frac{e\hbar}{2m_p c}$$

Q: What happens if we follow a measurement of  $S_z$  by a measurement of  $S_x$ ?



Each eigenstate of  $S_z$  is an equal superposition of  $S_x$  eigenstates.



Schematic of triple  
measurement.