

## Practice Problems for Physics 560A Midterm

Calculator and crib sheet (1-side 8.5" x 11") allowed. Show work for full credit.

### 1) Reciprocal lattice

Consider a three-dimensional Bravais lattice  $\mathbf{R} = n_1\mathbf{a}_1 + n_2\mathbf{a}_2 + n_3\mathbf{a}_3$ , where  $n_i \in \mathbb{Z}$  and  $\mathbf{a}_1$ ,  $\mathbf{a}_2$ , and  $\mathbf{a}_3$  are noncoplanar vectors.

a) Any function with the periodicity of the Bravais lattice may be expressed as a Fourier sum over a set of *reciprocal lattice vectors*. State the condition(s) which must be satisfied by a reciprocal lattice vector.

b) Consider an *orthorhombic* Bravais lattice, with fundamental translation vectors  $\mathbf{a}_1 = a\hat{x}$ ,  $\mathbf{a}_2 = b\hat{y}$ , and  $\mathbf{a}_3 = c\hat{z}$ , with  $a$ ,  $b$ , and  $c$  all different. What are the fundamental translation vectors of the reciprocal lattice?

c) What is the first Brillouin zone for the orthorhombic lattice in (b)?

### 2) Free-Electron Model

The heat capacity of a three-dimensional metal has the form  $C_V = AT$  at low temperatures. Give a detailed derivation of this temperature dependence using the free-electron model.

Useful integral:

$$\int_{-\infty}^{\infty} dx \frac{x^2}{\cosh^2 x/2} = 4\pi^2/3.$$

### 3) Zero-point fluctuations of a harmonic chain

Consider a one-dimensional Bravais lattice of atoms with lattice spacing  $a$ . The Hamiltonian is

$$H = \sum_{n=1}^L \left[ \frac{p_n^2}{2m} + \frac{C}{2} (x_n - x_{n-1})^2 \right].$$

Calculate the mean-squared deviation  $\langle 0|x_n^2|0\rangle$  of the  $n$ th atom from its equilibrium position at zero temperature.

Hint: the displacement operator for the  $n$ th atom may be expressed as

$$x_n = L^{-1/2} \sum_k \sqrt{\frac{\hbar}{2m\omega_k}} (a_k e^{ikna} + a_k^\dagger e^{-ikna}),$$

where  $\omega_k = \sqrt{4C/m} |\sin(ka/2)|$ . Also, in the limit  $L \rightarrow \infty$  (which you may utilize) the sum over  $k$  may be replaced by an integral,  $L^{-1} \sum_k \rightarrow (a/2\pi) \int dk$ . (In evaluating the sum over normal modes, the mode with  $k = 0$  should be omitted. This corresponds to displacements of the center of mass, and we are only interested in the displacement of the atom relative to a frame in which the center of mass of the crystal is fixed.)

What does your result imply about the possible existence of crystalline order in one dimension?