

Exercises for Physics 560A

Problem Set 8; Due November 26

1) Sharvin formula

The Sharvin formula for the electrical conductance of an extremely short contact of area A between two pieces of metal is

$$G \simeq \frac{2e^2}{h} \frac{k_F^2 A}{4\pi}.$$

Derive the Sharvin formula by considering the total current flowing through a hole of area A in a thin insulating barrier separating two free electron gases with different Fermi energies. Use purely macroscopic arguments. Hint: In a free electron gas, the number of electrons with energy between E and $E + dE$ traveling at an angle between θ and $\theta + d\theta$ with respect to a given axis is

$$\frac{\partial^2 n}{\partial E \partial \theta} dE d\theta = \frac{D(E)}{2} \sin \theta dE d\theta,$$

where $D(E)$ is the density of states.

2) Solution of Schrödinger equation inside a cylinder

Verify that the wavefunction

$$\psi_{kmn}(r, \phi, z) = e^{ikz + im\phi} J_m(\gamma_{mn}r/R)$$

is a solution of the free-particle Schrödinger equation in cylindrical coordinates, with energy eigenvalue

$$E = \frac{\hbar^2 \gamma_{mn}^2}{2mR^2} + \frac{\hbar^2 k^2}{2m}.$$

This wavefunction vanishes on the surface of a cylinder of radius R if γ_{mn} is the n th zero of J_m .

3) Conductance of a perfect cylindrical wire

Using the methods of Lecture 21 and the solution of problem 2, determine the electrical conductance of a perfect cylindrical wire of radius $k_F R = 6$.